

Master Theorem

Slides by Christopher M. Bourke
Instructor: Berthe Y. Choueiry

Spring 2006

Computer Science & Engineering 235
Introduction to Discrete Mathematics

When analyzing algorithms, recall that we only care about the *asymptotic behavior*.

Recursive algorithms are no different. Rather than *solve* exactly the recurrence relation associated with the cost of an algorithm, it is enough to give an asymptotic characterization.

The main tool for doing this is the *master theorem*.

Theorem (Master Theorem)

Let $T(n)$ be a monotonically increasing function that satisfies

$$\begin{aligned}T(n) &= aT\left(\frac{n}{b}\right) + f(n) \\T(1) &= c\end{aligned}$$

where $a \geq 1, b \geq 2, c > 0$. If $f(n) \in \Theta(n^d)$ where $d \geq 0$, then

$$T(n) = \begin{cases} \Theta(n^d) & \text{if } a < b^d \\ \Theta(n^d \log n) & \text{if } a = b^d \\ \Theta(n^{\log_b a}) & \text{if } a > b^d \end{cases}$$

You *cannot* use the Master Theorem if

- $T(n)$ is not monotone, ex: $T(n) = \sin n$
- $f(n)$ is not a polynomial, ex: $T(n) = 2T(\frac{n}{2}) + 2^n$
- b cannot be expressed as a constant, ex: $T(n) = T(\sqrt{n})$

Note here, that the Master Theorem does *not* solve a recurrence relation.

Does the base case remain a concern?

Master Theorem

Example 1

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = T\left(\frac{n}{2}\right) + \frac{1}{2}n^2 + n$. What are the parameters?

$a =$

$b =$

$d =$

Therefore which condition?

Master Theorem

Example 1

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = T\left(\frac{n}{2}\right) + \frac{1}{2}n^2 + n$. What are the parameters?

$$a = 1$$

$$b =$$

$$d =$$

Therefore which condition?

Master Theorem

Example 1

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = T\left(\frac{n}{2}\right) + \frac{1}{2}n^2 + n$. What are the parameters?

$$a = 1$$

$$b = 2$$

$$d =$$

Therefore which condition?

Master Theorem

Example 1

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = T\left(\frac{n}{2}\right) + \frac{1}{2}n^2 + n$. What are the parameters?

$$a = 1$$

$$b = 2$$

$$d = 2$$

Therefore which condition?

Master Theorem

Example 1

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = T\left(\frac{n}{2}\right) + \frac{1}{2}n^2 + n$. What are the parameters?

$$a = 1$$

$$b = 2$$

$$d = 2$$

Therefore which condition?

Since $1 < 2^2$, case 1 applies.

Master Theorem

Example 1

Let $T(n) = T\left(\frac{n}{2}\right) + \frac{1}{2}n^2 + n$. What are the parameters?

$$a = 1$$

$$b = 2$$

$$d = 2$$

Therefore which condition?

Since $1 < 2^2$, case 1 applies.

Thus we conclude that

$$T(n) \in \Theta(n^d) = \Theta(n^2)$$

Master Theorem

Example 2

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 2T\left(\frac{n}{4}\right) + \sqrt{n} + 42$. What are the parameters?

$a =$

$b =$

$d =$

Therefore which condition?

Master Theorem

Example 2

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 2T\left(\frac{n}{4}\right) + \sqrt{n} + 42$. What are the parameters?

$$a = 2$$

$$b =$$

$$d =$$

Therefore which condition?

Master Theorem

Example 2

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 2T\left(\frac{n}{4}\right) + \sqrt{n} + 42$. What are the parameters?

$$a = 2$$

$$b = 4$$

$$d =$$

Therefore which condition?

Master Theorem

Example 2

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 2T\left(\frac{n}{4}\right) + \sqrt{n} + 42$. What are the parameters?

$$a = 2$$

$$b = 4$$

$$d = \frac{1}{2}$$

Therefore which condition?

Master Theorem

Example 2

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 2T\left(\frac{n}{4}\right) + \sqrt{n} + 42$. What are the parameters?

$$a = 2$$

$$b = 4$$

$$d = \frac{1}{2}$$

Therefore which condition?

Since $2 = 4^{\frac{1}{2}}$, case 2 applies.

Master Theorem

Example 2

Let $T(n) = 2T\left(\frac{n}{4}\right) + \sqrt{n} + 42$. What are the parameters?

$$a = 2$$

$$b = 4$$

$$d = \frac{1}{2}$$

Therefore which condition?

Since $2 = 4^{\frac{1}{2}}$, case 2 applies.

Thus we conclude that

$$T(n) \in \Theta(n^d \log n) = \Theta(\sqrt{n} \log n)$$

Master Theorem

Example 3

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 3T\left(\frac{n}{2}\right) + \frac{3}{4}n + 1$. What are the parameters?

$a =$

$b =$

$d =$

Therefore which condition?

Master Theorem

Example 3

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 3T\left(\frac{n}{2}\right) + \frac{3}{4}n + 1$. What are the parameters?

$$a = 3$$

$$b =$$

$$d =$$

Therefore which condition?

Master Theorem

Example 3

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 3T\left(\frac{n}{2}\right) + \frac{3}{4}n + 1$. What are the parameters?

$$a = 3$$

$$b = 2$$

$$d =$$

Therefore which condition?

Master Theorem

Example 3

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 3T\left(\frac{n}{2}\right) + \frac{3}{4}n + 1$. What are the parameters?

$$a = 3$$

$$b = 2$$

$$d = 1$$

Therefore which condition?

Master Theorem

Example 3

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 3T\left(\frac{n}{2}\right) + \frac{3}{4}n + 1$. What are the parameters?

$$a = 3$$

$$b = 2$$

$$d = 1$$

Therefore which condition?

Since $3 > 2^1$, case 3 applies.

Master Theorem

Example 3

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 3T\left(\frac{n}{2}\right) + \frac{3}{4}n + 1$. What are the parameters?

$$a = 3$$

$$b = 2$$

$$d = 1$$

Therefore which condition?

Since $3 > 2^1$, case 3 applies. Thus we conclude that

$$T(n) \in \Theta(n^{\log_b a}) = \Theta(n^{\log_2 3})$$

Master Theorem

Example 3

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Let $T(n) = 3T\left(\frac{n}{2}\right) + \frac{3}{4}n + 1$. What are the parameters?

$$a = 3$$

$$b = 2$$

$$d = 1$$

Therefore which condition?

Since $3 > 2^1$, case 3 applies. Thus we conclude that

$$T(n) \in \Theta(n^{\log_b a}) = \Theta(n^{\log_2 3})$$

Note that $\log_2 3 \approx 1.5849\dots$. Can we say that
 $T(n) \in \Theta(n^{1.5849})$?

“Fourth” Condition

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Recall that we cannot use the Master Theorem if $f(n)$ (the non-recursive cost) is not polynomial.

There is a limited 4-th condition of the Master Theorem that allows us to consider polylogarithmic functions.

Corollary

If $f(n) \in \Theta(n^{\log_b a} \log^k n)$ for some $k \geq 0$ then

$$T(n) \in \Theta(n^{\log_b a} \log^{k+1} n)$$

This final condition is fairly limited and we present it merely for completeness.

“Fourth” Condition

Example

Master
Theorem

CSE235

Introduction

Pitfalls

Examples

4th Condition

Say that we have the following recurrence relation:

$$T(n) = 2T\left(\frac{n}{2}\right) + n \log n$$

Clearly, $a = 2, b = 2$ but $f(n)$ is not a polynomial. However,

$$f(n) \in \Theta(n \log n)$$

for $k = 1$, therefore, by the 4-th case of the Master Theorem we can say that

$$T(n) \in \Theta(n \log^2 n)$$